



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Fortunately, we have no rational use for flat roofs. Our cities are roomy, and our habits and their population will for many centuries keep them so. Our houses are low and our yards airy. I cannot conceive a single argument in favour either of the beauty or utility of terrace roofs in our country. Those that have them scarcely ever use them. The cold in winter and the heat in summer drive us from them. A beautiful prospect may justify the partial use of them, in particular situations, but neither architectural beauty, nor the general wants of our wintry climate call for their introduction.—To the southward beyond the reach of frost, however, the information contained in this paper may be highly useful.

No. LVII.

A general method of finding the roots of numeral equations to any degree of exactness; with the application of Logarithms to shorten the operation: by John Garnett of New Brunswick N. Jersey.

Read January 20th, 1809.

Suppose an equation, $ax + bx^2 + cx^3 + dx^4 + ex^5$ &c. = v , to find x .

RULE.

Find, by trial, any near root as x'

Then, by substitution, $ax' + bx'^2 + cx'^3 + dx'^4 + ex'^5$ &c. = v'

Multiply each term by the index of the power of x' , and divide by x' .

Let the products, $a + 2bx' + 3cx'^2 + 4dx'^3 + 5ex'^4$, &c. = A .

Multiply each term by the power of x' , and divide by $2x'$.

Let the products, $b + 3cx' + 6dx'^2 + 10ex'^3$, &c. = B .

Multiply each term by the power of x' , and divide by $3x'$.

Let the products, $c + 4dx' + 10ex'^2$, &c. = C .

Multiply each term again by its power of x' , and divide by $4x'$.

Let the products, $d + 5ex'$, &c. = D : and so on, continually, until all the powers of x' are destroyed; so that e , &c. = E .

Then will $Ax'' + Bx''^2 + Cx''^3 + Dx''^4 + Ex''^5$ &c. = $v - v'$, be a *New Equation* whose roots will all be less by x' , and the value, $v - v'$, less by v' than the roots of the original equation. And if the roots and value of this new equation be diminished in the same manner, by another near root, as x'' , and so on, continually, the root and value may become less than any assignable quantity, and the sum of all the near roots will be equal to x , the root of the original equation.

This rule will be found virtually the same as that given by Newton, Raphson, Jones, Simpson &c. and, if applied to the extraction of simple powers, will be found the same precisely as the one usually given in common arithmetic, thus for

EXAMPLE I.

$x' = 400$	Let $x^3 =$	99252847			
		64000000	$= x'^3$	$x' = 400$	
By the Rule,	$v - v'$	35252847	Resolvend.		
divisor		28800000	New Equation $x'' = 60$		
$3x'^2 = 480000 = A$		4320000	$480000 \times x''$		
$3x' = 1200 = B$		216000	$+ 1200 \times x''^2$		
$1 = 1 = C$			$+ 1 \times x''^3 = v - v'$		
By the Rule,		33336000	Subtrahend		
		01916847	Resolvend		
$A =$	divisor		New equation $x''' = 3$		
$480000 + 2400x'' + 3x''^2 = 634800.$		1904400	$634800 \times x'''$		
$B = 1200 + 3x = 1380.$		12420	$= 1380 \times x'''^2$		
$C = 1 = 1.$		27	$= 1 \times x'''^3$		
		1916847	Subtrahend.		

Whence $x = x' + x'' + x''' = 463$, the required Root.

This form will serve for all numeral equations, and with nearly the same labour; as for

EXAMPLE II.

$3x_3 + 2x^2 - 5x = v =$	242235792	near root $x' = 400$
$v' =$	192318000	$= 3x'^3 + 2x'^2 - 5x'$
By the Rule, $v - v' =$	49917792	Resolvend.
divisor	43247850	New equation $x'' = 30$
$9x'^2 + 4x' - 5 = 1441595 = A$	3241800	$= 1441595 \times x''$
$9x' + 2 = 3602 = B$	81000	$+ 3602 \times x''^2$
$3 = 3 = C$		$+ 3 \times x''^3 = v - v'$
v''	46570650	Subtrahend.
	3347142	Resolvend.
By the Rule,		$x''' = 2$
divisor	3331630	New Equation.
$9x''^2 + 7204x'' + 1441595 = 1665815$	15488	$1665815 \times x'''$
$9x + 3602 = 3872$	24	$+ 3872 \times x'''^2$
$3 = 3$		$+ 3 \times x'''^3 = v - v' - v''$
	3347142	Subtrahend.
Hence, $x = x' + x'' + x''' = 432.$	000000	

By dividing the original equation by $x-432=0$, the other roots may be found, but in this case they are imaginary. But, instead of thus approximating to a root by single figures, we can (after the root has been sufficiently diminished) find, by a Table of *Logarithms*, as many places of figures at one operation, as there are places of figures in the logarithms; as in the following

EXAMPLE III.

Suppose $x^3 - 2x = 5$. near Root $x' = 2$
 then $x'^3 - 2x' = 4$

By the Rule, divisor. Resolvend 1

$$\left. \begin{array}{r} 3x'^2 - 2 = 10 \\ 3x' = 6 \\ 1 = 1 \end{array} \right\}$$

near Root $x = .09$

gives $10x'' + 6x''^2 + x''^3 = 0.949329$ Subtrahend.

Resolvend $0.050671 = v$.

This being now sufficiently reduced we proceed thus:—

By the Rule, New Equation.

$$\left. \begin{array}{r} 3x''^2 + 12x'' + 10 = 11,1043 \text{ (A)} \\ 3x'' + 6 = 6,27 = B \\ 1 = 1 = C \end{array} \right\} \begin{array}{l} 11,1043x''' + 6,27x'''^2 + x'''^3 = 0,050671 = V \text{ and by} \\ \text{reversion of series, } x''' = \frac{V}{A} - \frac{B}{A^2}V^2 + \&c. \end{array}$$

Then, by logarithms,

$V = ,050671$	Log. v	8.704760
$A = 11,1043$	Log. a	1.045492
$\frac{V}{A} = ,00456319$	Log. d	7.659268
$\frac{V}{A^2}$	c	6.613776
$\frac{B}{A^2}$	b	0.797268
$\frac{B}{A^3}V^2 = ,00001176$		5.070312
$\frac{V}{A} - \frac{B}{A^2}V^2 = x''' = ,00455143$		

whence $x = 2.09455143$

And if ,004551 be put for x''' , in the above new equation, the value and root would be again reduced, so that we should obtain $x'''' = ,0000004815424$, and consequently the root $x = 2,0945514815424$, true to the last figure.